

t33_xreal_1 (TMa-
toH4HSD3PEgcjvjyQLvu3HtmYPwMWCSU)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (((\neg v3_xxreal_0 X0) \wedge (v1_xreal_0 X0)) \wedge ((\neg v3_xxreal_0 X1) \wedge (v1_xreal_0 X1))) \Rightarrow (\neg v3_xxreal_0 (k2_xcmplx_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow ((v3_xxreal_0 X0) \Leftrightarrow (\neg r1_xxreal_0 k6_numbers X0)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (4)$$

Theorem 1

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 k6_numbers X1)) \Rightarrow (r1_xxreal_0 k6_numbers (k2_xcmplx_0 X0 X1))))$$