

t33_xxreal_0
(TMTFscjSUL21P6B13hpj9tEdQLpUXgLwCHg)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k3_xxreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & (r1_xxreal_0 X0 X1) \Rightarrow (k3_xxreal_0 X0 X1 = X0)) \wedge ((\neg r1_xxreal_0 X0 \\ & X1) \Rightarrow (k3_xxreal_0 X0 X1 = X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (\\ & k3_xxreal_0 X0 X1 = k3_xxreal_0 X1 X0) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (k3_xxreal_0 (k3_xxreal_0 X0 X1) X2 = k3_xxreal_0 \\ & X0 (k3_xxreal_0 X1 X2)))) \end{aligned}$$