

t33_xxreal_1
(TMXD9JR6qBK5sgrhKgxqSShRwqknyzxPLLk)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\neg \\ & (\neg r1_xxreal_0 X1 X0) \wedge (\forall X2. (v1_xxreal_0 X2) \Rightarrow (\neg(\neg r1_xxreal_0 \\ & X2 X0) \wedge (\neg r1_xxreal_0 X1 X2)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\neg \\ & (\neg r1_xxreal_0 X1 X0) \wedge (v1_xboole_0 (k4_xxreal_1 X0 X1)))) \end{aligned}$$