

t34_asympt_0

(TMQDy3cgC7eT8GE2RjGHc86Bix2ng7bW5ru)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v4_asympt_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_comseq_2 : \iota \Rightarrow o$ be given. Let $k52_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_seq_2 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k6_asympt_0 : \iota \Rightarrow \iota$ be given. Let $k8_asympt_0 : \iota \Rightarrow \iota$ be given. Let $v2_asympt_0 : \iota \Rightarrow o$ be given. Let $k7_asympt_0 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_asympt_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & ((v2_asympt_0 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k1_numbers)))))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 k5_numbers k1_numbers) \wedge ((v2_asympt_0 X1) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((\\ & X0 \in k7_asympt_0 X1) \Leftrightarrow (X1 \in k6_asympt_0 X0)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & ((v4_asympt_0 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k1_numbers)))))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 k5_numbers k1_numbers) \wedge ((v4_asympt_0 X1) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((\\ & (v2_comseq_2 (k52_valued_1 k5_numbers k1_numbers k1_numbers \\ & X0 X1) \wedge (k2_seq_2 (k52_valued_1 k5_numbers k1_numbers k1_numbers \\ & X0 X1) = k6_numbers)) \Rightarrow ((X0 \in k6_asympt_0 X1) \wedge (\neg X1 \in k6_asympt_0 \\ & X0)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. \\ & (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & ((v2_asympt_0 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k1_numbers)))))) \Rightarrow (k8_asympt_0 X0 = k3_xboole_0 (k6_asympt_0 \\ & X0) (k7_asympt_0 X0)) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & k1_numbers))) \Rightarrow (((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers \\ & k1_numbers) \wedge (v4_asympt_0 X0))) \Rightarrow ((v1_funct_1 X0) \wedge ((v1_funct_2 \\ & X0 k5_numbers k1_numbers) \wedge ((v2_asympt_0 X0) \wedge (v5_asympt_0 X0)))))) \end{aligned} \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & ((v4_asympt_0 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers k1_numbers)))))) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 k5_numbers k1_numbers) \wedge ((v4_asympt_0 X1) \wedge (m1_subset_1 \\ & X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow ((\\ & (v2_comseq_2 (k52_valued_1 k5_numbers k1_numbers k1_numbers \\ & X0 X1)) \wedge (k2_seq_2 (k52_valued_1 k5_numbers k1_numbers k1_numbers \\ & X0 X1) = k6_numbers)) \Rightarrow ((X0 \in k6_asympt_0 X1) \wedge (\neg X0 \in k8_asympt_0 \\ & X1)))) \end{aligned}$$