

t34\_complex2  
(TMGbBe77njxFe8EPCY8Yn26NxdtHfL1n8wf)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_numbers : \iota$  be given. Let  $k1\_complex2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k15\_complex1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow (k15\_complex1 (k3\_xcmplx\_0 X0 X1) = k9\_complex1 (k15\_complex1 X0) (k15\_complex1 X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k2\_numbers) \wedge (m1\_subset\_1 X1 k2\_numbers)) \Rightarrow (k9\_complex1 X0 X1 = k3\_xcmplx\_0 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k15\_complex1 (k15\_complex1 X0) = X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (m1\_subset\_1 (k15\_complex1 X0) k2\_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcmplx\_0 X1) \Rightarrow (k1\_complex2 X0 X1 = k3\_xcmplx\_0 X0 (k15\_complex1 X1))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k2\_numbers) \wedge (m1\_subset\_1 X1 k2\_numbers)) \Rightarrow (k9\_complex1 X0 X1 = k9\_complex1 X1 X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k3\_xcmplx\_0 X0 X1 = k3\_xcmplx\_0 X1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k2\_numbers) \Rightarrow (v1\_xcmplx\_0 X0) \quad (8)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 k2\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 X1 k2\_numbers) \Rightarrow (k1\_complex2 X0 X1 = k15\_complex1 (k1\_complex2 X1 X0)))$$