

### t34\_comseq\_3

(TMSvjHeFRBdu7LugRdLWfd2un645ReX9V77)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_valued\_0 : \iota \Rightarrow o$  be given. Let  $k3\_series\_1 : \iota \Rightarrow \iota$  be given. Let  $k55\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k3\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k8\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_membered : \iota \Rightarrow o$  be given. Let  $k54\_valued\_1 : \iota \Rightarrow \iota$  be given. Let  $k17\_complex1 : \iota \Rightarrow \iota$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))) \Rightarrow \\ & ((\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow (r1\_xxreal\_0 \\ & k6\_numbers (k3\_funct\_2 k5\_numbers k1\_numbers X0 X1))) \Rightarrow (v7\_valued\_0 \\ & (k3\_series\_1 X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 \\ & X1 k5\_numbers X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & X0)))) \wedge (v7\_ordinal1 X2)) \Rightarrow (k8\_nat\_1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \end{aligned} \tag{3}$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered\ X1)\wedge((v1\_funct\_1\ X2)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1))))))\Rightarrow(k55\_valued\_1\ X0\ X1\ X2 = k54\_valued\_1\ X2) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1\_xboole\_0\ X0)\wedge(((v1\_funct\_1\ X2)\wedge((v1\_funct\_2\ X2\ X0\ X1)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1))))))\wedge(m1\_subset\_1\ X3\ X0)))\Rightarrow(k3\_funct\_2\ X0\ X1\ X2\ X3 = k1\_funct\_1\ X2\ X3) \quad (6)$$

Assume the following.

$$\forall X0.((v1\_funct\_1\ X0)\wedge((v1\_funct\_2\ X0\ k5\_numbers\ k2\_numbers)\wedge(m1\_subset\_1\ X0\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ k5\_numbers\ k2\_numbers))))))\Rightarrow(\forall X1.(m2\_subset\_1\ X1\ k1\_numbers\ k5\_numbers)\Rightarrow((k17\_complex1\ (k8\_nat\_1\ k2\_numbers\ X0\ X1) = k8\_nat\_1\ k1\_numbers\ (k55\_valued\_1\ k5\_numbers\ k2\_numbers\ X0)\ X1)\wedge(r1\_xxreal\_0\ k6\_numbers\ (k8\_nat\_1\ k1\_numbers\ (k55\_valued\_1\ k5\_numbers\ k2\_numbers\ X0)\ X1)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v1\_xboole\_0\ X1)\wedge(v1\_membered\ X1))\wedge((v1\_funct\_1\ X2)\wedge((v1\_funct\_2\ X2\ X0\ X1)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1))))))\Rightarrow((v1\_funct\_1\ (k54\_valued\_1\ X2))\wedge(v1\_partfun1\ (k54\_valued\_1\ X2)\ X0)) \quad (8)$$

Assume the following.

$$(\neg v1\_xboole\_0\ k4\_ordinal1)\wedge(v3\_ordinal1\ k4\_ordinal1) \quad (9)$$

Assume the following.

$$\neg v1\_xboole\_0\ k2\_numbers \quad (10)$$

Assume the following.

$$\neg v1\_xboole\_0\ k1\_numbers \quad (11)$$

Assume the following.

$$v1\_membered\ k2\_numbers \quad (12)$$

Assume the following.

$$m1\_subset\_1\ k5\_numbers\ (k1\_zfmisc\_1\ k1\_numbers) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered\ X1)\wedge((v1\_funct\_1\ X2)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1))))))\Rightarrow((v1\_funct\_1\ (k55\_valued\_1\ X0\ X1\ X2))\wedge(m1\_subset\_1\ (k55\_valued\_1\ X0\ X1\ X2)\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ k1\_numbers)))) \quad (14)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v1\_partfun1 X2 X0) \Rightarrow (v1\_funct\_2 X2 X0 X1)) \quad (16)$$

**Theorem 1**

$$\forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k2\_numbers) \wedge (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k2\_numbers)))))) \Rightarrow (v7\_valued\_0 (k3\_series\_1 (k55\_valued\_1 k5\_numbers k2\_numbers X0)))$$