

t34_fib_num3 (TMWDu- JDThqPuoeo7nPzSEZGjnMARtjGQgTk)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_fib_num3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(v7_ordinal1\ X1) \Rightarrow ((\\ k3_fib_num3\ X0\ X1\ k6_numbers = X0) \wedge ((k3_fib_num3\ X0\ X1\ np_1 = X1) \wedge \\ (\forall X2.(v7_ordinal1\ X2) \Rightarrow (k3_fib_num3\ X0\ X1\ (k2_nat_1\ (k1_nat_1 \\ X2\ np_1)\ np_1) = k2_nat_1\ (k3_fib_num3\ X0\ X1\ X2)\ (k3_fib_num3\ X0 \\ X1\ (k1_nat_1\ X2\ np_1))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v1_xcmplx_0\ X0) \wedge ((v1_xcmplx_0 \\ X1) \wedge (v1_xcmplx_0\ X2))) \Rightarrow (k2_xcmplx_0\ (k2_xcmplx_0\ X0\ X1)\ X2 = k2_xcmplx_0 \\ X0\ (k2_xcmplx_0\ X1\ X2)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0\ np_2) \wedge (m2_subset_1\ np_2\ k1_numbers\ k5_numbers)) \wedge \\ ((m1_subset_1\ np_2\ k5_numbers) \wedge (m1_subset_1\ np_2\ k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} ((v2_xxreal_0\ np_1) \wedge (m2_subset_1\ np_1\ k1_numbers\ k5_numbers)) \wedge \\ ((m1_subset_1\ np_1\ k5_numbers) \wedge (m1_subset_1\ np_1\ k1_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k2_xcmplx_0\ np_1\ np_1 = np_2 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow(k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow(k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \quad (11)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xcmplx_0 X0) \quad (14)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow(k2_nat_1 (k3_fib_num3 X0 X1 X2) (k3_fib_num3 X0 X1 (k1_nat_1 X2 np_1)) = k3_fib_num3 X0 X1 (k1_nat_1 X2 np_2))))$$