

t34_int_1

(TMXLw8Bj4NnSNuodGGZJnaWbWR4HNqxSMcC)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_int_1 : \iota \Rightarrow \iota$ be given. Let $k2_int_1 : \iota \Rightarrow \iota$ be given. Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k3_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((\neg r1_xxreal_0 (k2_int_1 X0) (k5_real_1 X0 np_1)) \wedge (\neg r1_xxreal_0 (k3_real_1 (k2_int_1 X0) np_1) X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((\neg(\neg r1_xxreal_0 (k2_int_1 X0) X0) \wedge (v1_int_1 X0)) \wedge (\neg(\neg v1_int_1 X0) \wedge (r1_xxreal_0 (k2_int_1 X0) X0))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_int_1 (k2_int_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow ((X1 = k1_int_1 X0) \Leftrightarrow ((r1_xxreal_0 X1 X0) \wedge (\neg r1_xxreal_0 X1 (k5_real_1 X0 np_1)))))) \quad (4)$$

Theorem 1 $\forall X0.(v1_xreal_0 X0) \Rightarrow ((k1_int_1 X0 = k2_int_1 X0) \Leftrightarrow (v1_int_1 X0)).$