

t34_jgraph_1

(TMYeBN1u58STkdUhXa3jKSfjWvss7rEWRJF)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k9_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k5_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (k5_algstr_0 (k15_euclid np_2) X0 X1 = k19_euclid (k9_real_1 (\\ & k17_euclid X0) (k17_euclid X1)) (k9_real_1 (k18_euclid X0) (k18_euclid \\ & X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((r1_xreal_0 (k18_complex1 (k17_euclid X0)) (k12_euclid X0)) \wedge \\ & (r1_xreal_0 (k18_complex1 (k18_euclid X0)) (k12_euclid X0))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k9_real_1 X0 X1) k1_numbers) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (m1_subset_1 (k19_euclid X0 X1) (u1_struct_0 (k15_euclid np_2))) \tag{5}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (m1_subset_1 (k18_euclid X0) k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow (m1_subset_1 (k17_euclid X0) k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((r1_xxreal_0 (k18_complex1 (k9_real_1 (k17_euclid X0) (k17_euclid \\ & X1))) (k12_euclid (k5_algstr_0 (k15_euclid np_2) X0 X1))) \wedge (r1_xxreal_0 \\ & (k18_complex1 (k9_real_1 (k18_euclid X0) (k18_euclid X1))) (k12_euclid \\ & (k5_algstr_0 (k15_euclid np_2) X0 X1)))))) \end{aligned}$$