

t34\_newton

(TMHX6v1dDtFVv6SEP1dXFFM3Cvkcqi45qJC)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 np\_1 X0) \Rightarrow (r1\_xxreal\_0 X1 (k3\_xcmplx\_0 X1 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0) \wedge (v7\_ordinal1 X1)) \Rightarrow (v7\_ordinal1 (k3\_xcmplx\_0 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_xreal\_0 X0) \quad (4)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2.(v7\_ordinal1 X2) \Rightarrow (((r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 (k3\_xcmplx\_0 X2 X0) X1)) \Rightarrow (r1\_xxreal\_0 X2 X1))))$$