

t34\_procal\_1  
(TMKd7QJKkF7g5MrDMbR5KZgjLhphoPtffsA)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_cqc\_the1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (k8\_cqc\_lang X0 X1 (k9\_cqc\_lang X0 X2 X1) \in \\ k4\_cqc\_the1 X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (((k8\_cqc\_lang X0 X1 X2 \in k4\_cqc\_the1 X0) \wedge \\ (k8\_cqc\_lang X0 X2 X3 \in k4\_cqc\_the1 X0)) \Rightarrow (k8\_cqc\_lang X0 X1 X3 \in k4\_cqc\_the1 \\ X0)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow ((k8\_cqc\_lang X0 X1 X2 \in k4\_cqc\_the1 X0) \Rightarrow \\ (k8\_cqc\_lang X0 (k8\_cqc\_lang X0 X2 X3) (k8\_cqc\_lang X0 X1 X3) \in k4\_cqc\_the1 \\ X0)))))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k3\_cqc\_lang X0)) \quad (5)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k9\_qc\_lang1 X0)) \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1\_qc\_lang1 X0) \wedge ((m1\_subset\_1 \\ & X1 (k3\_cqc\_lang X0)) \wedge (m1\_subset\_1 X2 (k3\_cqc\_lang X0)))) \Rightarrow (m2\_subset\_1 \\ & (k9\_cqc\_lang X0 X1 X2) (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((m1\_qc\_lang1 X0) \wedge ((m1\_subset\_1 \\ & X1 (k3\_cqc\_lang X0)) \wedge (m1\_subset\_1 X2 (k3\_cqc\_lang X0)))) \Rightarrow (m2\_subset\_1 \\ & (k8\_cqc\_lang X0 X1 X2) (k9\_qc\_lang1 X0) (k3\_cqc\_lang X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (m1\_subset\_1 (k3\_cqc\_lang X0) (k1\_zfmisc\_1 (k9\_qc\_lang1 X0))) \quad (9)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 \\ & X0) (k3\_cqc\_lang X0)) \Rightarrow (k8\_cqc\_lang X0 (k8\_cqc\_lang X0 (k9\_cqc\_lang \\ & X0 X1 X2) X3) (k9\_cqc\_lang X0 (k8\_cqc\_lang X0 X1 X3) (k8\_cqc\_lang \\ & X0 X2 X3)) \in k4\_cqc\_the1 X0)))) \end{aligned}$$