

t34_rfunct_1
(TMYd7XrhxaaZvhrb3nsy83Rd3H3ENNbsiA9)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k18_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_rfunct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_rfunct_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 \\ & \quad X1)))) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_valued_0 \\ & \quad X2)))) \Rightarrow (k18_valued_1 (k18_valued_1 X0 X1) X2 = k18_valued_1 X0 (\\ & \quad k18_valued_1 X1 X2))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 \\ & \quad X1)))) \Rightarrow (k1_rfunct_1 X0 X1 = k18_valued_1 X0 (k4_rfunct_1 X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 \\ & \quad X1)))) \Rightarrow (k4_rfunct_1 (k18_valued_1 X0 X1) = k18_valued_1 (k4_rfunct_1 \\ & \quad X0) (k4_rfunct_1 X1))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & \quad ((v1_relat_1 (k4_rfunct_1 X0)) \wedge ((v1_funct_1 (k4_rfunct_1 X0)) \wedge \\ & \quad (v1_valued_0 (k4_rfunct_1 X0)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1.(((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 \\ & \quad X0))) \wedge ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 X1)))) \Rightarrow \\ & \quad ((v1_relat_1 (k18_valued_1 X0 X1)) \wedge ((v1_funct_1 (k18_valued_1 \\ & \quad X0 X1)) \wedge (v1_valued_0 (k18_valued_1 X0 X1)))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 \\ X0)))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_valued_0 X1))))\Rightarrow \quad (6) \\ (k18_valued_1 X0 X1 = k18_valued_1 X1 X0) \end{aligned}$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow \\ (\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_valued_0 \\ X1))))\Rightarrow(\forall X2.((v1_relat_1 X2)\wedge((v1_funct_1 X2)\wedge(v1_valued_0 \\ X2))))\Rightarrow(\forall X3.((v1_relat_1 X3)\wedge((v1_funct_1 X3)\wedge(v1_valued_0 \\ X3))))\Rightarrow(k18_valued_1 (k1_rfunct_1 X0 X1) (k1_rfunct_1 X2 X3) = k1_rfunct_1 \\ (k18_valued_1 X0 X2) (k18_valued_1 X1 X3)))) \end{aligned}$$