

t34_rvsum_1 (TMZRPxJQk-
WgF1EbQvAErjMrhntD4W2c5RZe)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k8_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $k4_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (k8_rvsum_1 X0 X1 = k4_rvsum_1 X0 (k6_rvsum_1 X1))) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (k6_rvsum_1 X0 = k30_valued_1 X0) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (k6_rvsum_1 (k6_rvsum_1 X0) = X0) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_valued_0 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow ((v1_relat_1 (k30_valued_1 X0)) \wedge ((v1_funct_1 (k30_valued_1 X0)) \wedge ((v1_valued_0 (k30_valued_1 X0)) \wedge (v1_finseq_1 (k30_valued_1 X0)))))) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v3_valued_0 X0))) \Rightarrow ((v1_relat_1 (k30_valued_1 X0)) \wedge ((v1_funct_1 (k30_valued_1 X0)) \wedge ((v1_valued_0 (k30_valued_1 X0)) \wedge (v3_valued_0 (k30_valued_1 X0))))) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \Rightarrow ((v1_relat_1 X0) \wedge (v1_valued_0 X0)) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ & X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 \\ & X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (k8_rvsum_1 X0 (k6_rvsum_1 \\ & X1) = k4_rvsum_1 X0 X1)) \end{aligned}$$