

t34_tsp_2

(TMUMPW9wvS8qiVGphmmuUTu2PJX9wPcS8Nf)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v2_tsp_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tsp_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_tex_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 \\ X0))) \Rightarrow ((v3_pre_topc X1 X0) \Rightarrow (k3_tex_4 X0 X1 = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_pre_topc X1 X0) \Rightarrow \\ (m1_subset_1 (u1_struct_0 X1) (k1_zfmisc_1 (u1_struct_0 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 \\ X0)) \Rightarrow (k8_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_pre_topc X1 X0) \Rightarrow \\ (l1_pre_topc X1)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(m1_subset_1 (k7_relset_1 X0 X1 X2 X3) (k1_zfmisc_1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((\neg v2_struct_0 X0)\wedge((v2_pre_topc X0)\wedge(l1_pre_topc X0)))\wedge((\neg v2_struct_0 X1)\wedge((v2_tsp_2 X1 X0)\wedge(m1_pre_topc X1 X0))))\Rightarrow((v1_funct_1 (k1_tsp_2 X0 X1))\wedge((v1_funct_2 (k1_tsp_2 X0 X1) (u1_struct_0 X0) (u1_struct_0 X1))\wedge((v5_pre_topc (k1_tsp_2 X0 X1) X0 X1)\wedge(m1_subset_1 (k1_tsp_2 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X1)))))))) \quad (7)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0)\Rightarrow(\forall X1.(l1_pre_topc X1)\Rightarrow((m1_pre_topc X1 X0)\Leftrightarrow((r1_tarski (u1_struct_0 X1) (u1_struct_0 X0))\wedge(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X1)))\Rightarrow((v3_pre_topc X2 X1)\Leftrightarrow(\exists X3.(m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0)))\wedge((v3_pre_topc X3 X0)\wedge(X2 = k8_subset_1 (u1_struct_0 X0) X3 (u1_struct_0 X1)))))))))) \quad (8)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0)\wedge((v2_pre_topc X0)\wedge(l1_pre_topc X0)))\Rightarrow(\forall X1.((\neg v2_struct_0 X1)\wedge((v2_tsp_2 X1 X0)\wedge(m1_pre_topc X1 X0)))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 X2 (u1_struct_0 X0) (u1_struct_0 X1))\wedge((v5_pre_topc X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X1))))))))\Rightarrow((X2 = k1_tsp_2 X0 X1)\Leftrightarrow(\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0)))\Rightarrow((X3 = u1_struct_0 X1)\Rightarrow(\forall X4.(m1_subset_1 X4 (k1_zfmisc_1 (u1_struct_0 X0)))\Rightarrow(k9_subset_1 (u1_struct_0 X0) X3 (k3_tex_4 X0 X4) = k7_relset_1 (u1_struct_0 X0) (u1_struct_0 X1) X2 X4)))))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (10)$$

Theorem 1

$$\forall X0.((\neg v2_struct_0 X0)\wedge((v2_pre_topc X0)\wedge(l1_pre_topc X0)))\Rightarrow(\forall X1.((\neg v2_struct_0 X1)\wedge((v2_tsp_2 X1 X0)\wedge(m1_pre_topc X1 X0)))\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0)))\Rightarrow((v3_pre_topc X2 X0)\Rightarrow(v3_pre_topc (k7_relset_1 (u1_struct_0 X0) (u1_struct_0 X1) (k1_tsp_2 X0 X1) X2) X1))))$$