

t35_funct_2

(TMTZs29FgMX6FjfBpUPwJjQmZDNY9aMLqAZ)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. (X4 \in k9_xtuple_0 X0) \wedge ((X4 \in X1) \wedge (X3 = k1_funct_1 X0 X4)))))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X3)\wedge \\ & ((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1))))\Rightarrow((X1\neq k1_xboole_0)\Rightarrow(\forall X4.(\exists X5.(X5 \in X0)\wedge \\ & ((X5 \in X2)\wedge(X4 = k1_funct_1 X3 X5)))\Rightarrow(X4 \in k7_relset_1 X0 X1 X3 X2))) \end{aligned}$$