

# t35\_funct\_4 (TM- Fzd7agGHMrZqARZH4DvCFi14ysE9ZFmeo)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_xboole\_0 (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \Rightarrow (r1\_partfun1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_partfun1 X0 X1) \Leftrightarrow (k1\_funct\_4 X0 X1 = k1\_funct\_4 X1 X0))) \quad (2)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_xboole\_0 (k9\_xtuple\_0 X0) (k9\_xtuple\_0 X1)) \Rightarrow (k1\_funct\_4 X0 X1 = k1\_funct\_4 X1 X0)))$$