

t35_jordan5d

(TMUMmsU9xiR1tmopdoSX2M5fRwhFGZKZcE1)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_6 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_topreal1 : \iota \Rightarrow o$ be given. Let $v2_topreal1 : \iota \Rightarrow o$ be given. Let $v1_goboard5 : \iota \Rightarrow o$ be given. Let $v2_goboard5 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_goboard2 : \iota \Rightarrow \iota$ be given. Let $k1_matrix_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k3_matrix_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_matrix_1 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_4 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. ((\neg v1_xboole_0 X0) \wedge (m2_finseq_1 X0 (u1_struct_0 (\\
 & \quad k15_euclid np_2)))) \Rightarrow (\forall X1. (m2_subset_1 X1 k1_numbers \\
 & \quad k5_numbers) \Rightarrow (\forall X2. (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow \\
 & \quad (\neg (r1_xxreal_0 np_1 X1) \wedge ((r1_xxreal_0 X1 (k3_finseq_1 (k2_goboard2 \\
 & \quad X0))) \wedge ((r1_xxreal_0 np_1 X2) \wedge ((r1_xxreal_0 X2 (k1_matrix_1 \\
 & \quad (k2_goboard2 X0))) \wedge (\forall X3. (m2_subset_1 X3 k1_numbers k5_numbers) \Rightarrow \\
 & \quad (\neg (X3 \in k4_finseq_1 X0) \wedge ((k4_tarski X1 X2 \in k2_matrix_1 (k2_goboard2 \\
 & \quad X0)) \wedge (k17_euclid (k7_partfun1 (u1_struct_0 (k15_euclid np_2)) \\
 & \quad X0 X3) = k17_euclid (k3_matrix_1 (u1_struct_0 (k15_euclid np_2)) \\
 & \quad (k2_goboard2 X0) X1 X2))))))))))
 \end{aligned}
 \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v3_funct_1 X0) \wedge ((\neg v1_xboole_0 X0) \wedge ((v1_finseq_6 \\ & X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 \\ & X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge (m2_finseq_1 X0 (u1_struct_0 \\ & (k15_euclid np_2)))))))))) \Rightarrow (\neg r1_xxreal_0 (k3_finseq_1 X0) \\ & np_4) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (((X1 \in k4_finseq_1 \\ & X0) \wedge (r1_xxreal_0 np_2 (k3_finseq_1 X0))) \Rightarrow (k7_partfun1 (u1_struct_0 \\ & (k15_euclid np_2)) X0 X1 \in k3_topreal1 np_2 X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_4) \wedge (m2_subset_1 np_4 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_4 k5_numbers) \wedge (m1_subset_1 np_4 k1_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\neg r1_xxreal_0 np_4 np_2 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (10)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(m1_subset_1 (k7_partfun1 X0 X1 X2) X0) \quad (14)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (15)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow((r1_xxreal_0 X0 X1)\vee(r1_xxreal_0 X1 X0)) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (19)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (20)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0)\wedge((\neg v3_funct_1 X0)\wedge((v1_finseq_6 X0 (u1_struct_0 (k15_euclid np_2)))\wedge((v1_topreal1 X0)\wedge((v2_topreal1 X0)\wedge((v1_goboard5 X0)\wedge((v2_goboard5 X0)\wedge(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))))))))))\Rightarrow(\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers)\Rightarrow(\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers)\Rightarrow(\neg(r1_xxreal_0 np_1 X1)\wedge((r1_xxreal_0 X1 (k3_finseq_1 (k2_goboard2 X0))\wedge((r1_xxreal_0 np_1 X2)\wedge((r1_xxreal_0 X2 (k1_matrix_1 (k2_goboard2 X0))\wedge(\forall X3.(m1_subset_1 X3 (u1_struct_0 (k15_euclid np_2)))\Rightarrow(\neg(k17_euclid X3 = k17_euclid (k3_matrix_1 (u1_struct_0 (k15_euclid np_2)) (k2_goboard2 X0) X1 X2))\wedge(X3 \in k3_topreal1 np_2 X0))))))))))$$