

t35\_series\_1 (TMT-  
nRh6AnHd8guhjJhmKVWcGaQkKtCxRRhv)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_series\_1 : \iota \Rightarrow o$  be given. Let  $v1\_series\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0. (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ k1\_numbers))) \Rightarrow (((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers \\ k1\_numbers) \wedge (v2\_series\_1 X0))) \Rightarrow ((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 \\ X0 k5\_numbers k1\_numbers) \wedge (v1\_series\_1 X0)))) \end{aligned} \quad (1)$$

**Theorem 1**

$$\begin{aligned} \forall X0. ((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ ((v2\_series\_1 X0) \Rightarrow (v1\_series\_1 X0)) \end{aligned}$$