

t368_xxreal_1 (TMQdcd-
FCXRKf4GVkN4hNueeAuVBzCMbQPv5)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k6_subset_1 (k1_xxreal_1 \\ & X0 k1_xxreal_0) (k1_xxreal_1 X2 X1) = k2_xboole_0 (k2_xxreal_1 \\ & X0 X2) (k3_xxreal_1 X1 k1_xxreal_0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k1_xxreal_1 X0 X0 = k1_tarski X0) \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((\\ & r1_xxreal_0 X0 X1) \Rightarrow (k6_subset_1 (k1_xxreal_1 X0 k1_xxreal_0) \\ & (k1_tarski X1) = k2_xboole_0 (k2_xxreal_1 X0 X1) (k3_xxreal_1 X1 \\ & k1_xxreal_0)))))) \end{aligned}$$