

## t36\_bvfunc25

(TMH6oC8K2q1UQP4sf96tVL2Hfhrdw87bGhZ)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\ (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge \\ ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ (k1\_bvfunc\_1 X0 (k9\_bvfunc\_1 X0 X1 X2)) (k2\_bvfunc\_1 X0 X1 (k1\_bvfunc\_1 \\ X0 X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. \forall X3. (((v1\_funct\_1 X2) \wedge \\ ((v1\_funct\_2 X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ X0 X1)))))) \wedge ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\ X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow ((r2\_funct\_2 X0 X1 X2 \\ X3) \Rightarrow (r2\_funct\_2 X0 X1 X3 X2)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 X1) \wedge ( \\ (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (k1\_bvfunc\_1 X0 (k1\_bvfunc\_1 \\ X0 X1) = X1) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 \\ & X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k9\_bvfunc\_1 X0 X1 X2)) \wedge ( \\ & (v1\_funct\_2 (k9\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ & (k9\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 \\ & X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k2\_bvfunc\_1 X0 X1 X2)) \wedge ( \\ & (v1\_funct\_2 (k2\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ & (k2\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k1\_bvfunc\_1 \\ & X0 X1)) \wedge ((v1\_funct\_2 (k1\_bvfunc\_1 X0 X1) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ & (k1\_bvfunc\_1 X0 X1) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (6)$$

**Theorem 1**

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge ( \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & (k2\_bvfunc\_1 X0 X1 X2) (k1\_bvfunc\_1 X0 (k9\_bvfunc\_1 X0 X1 (k1\_bvfunc\_1 \\ & X0 X2)))))) \end{aligned}$$