

t36_funct_7 (TMVcsEUjMbQBJgMAGx- EcwNssQzuV8yftPgk)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X2 = X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & k9_xtuple_0 (k2_funct_7 X0 X2 X1) = k9_xtuple_0 X0) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((m1_finseq_1 X1 \\ & X0) \wedge ((m1_subset_1 X2 k5_numbers) \wedge (m1_subset_1 X3 X0))) \Rightarrow (k3_funct_7 \\ & X0 X1 X2 X3 = k2_funct_7 X1 X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (\\ & (v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_finseq_1 X1 X0)\wedge((m1_subset_1 X2 k5_numbers)\wedge(m1_subset_1 X3 X0)))\Rightarrow(m2_finseq_1 (k3_funct_7 X0 X1 X2 X3) X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (9)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 X0)\Rightarrow(\forall X3.(m1_subset_1 X3 k5_numbers)\Rightarrow((X3 \in k4_finseq_1 X1)\Rightarrow(k7_partfun1 X0 (k3_funct_7 X0 X1 X3 X2) X3 = X2))))))$$