

t36_fuzzy_1

(TMQMT1JBwqDYfVCKw5mRxW5t4TyMTi8Ayzk)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_fuzzy_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_fuzzy_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v5_relat_1 X1 (k1_rcomp_1 \\ & k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ & (\forall X2. ((v5_relat_1 X2 (k1_rcomp_1 k6_numbers np_1)) \wedge (\\ & (v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 k1_numbers) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow ((r1_fuzzy_1 \\ & X1 X2) \Rightarrow (r1_fuzzy_1 (k3_fuzzy_1 X0 X2) (k3_fuzzy_1 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v5_relat_1 X1 (k1_rcomp_1 \\ & k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ & (k3_fuzzy_1 X0 (k3_fuzzy_1 X0 X1) = X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v5_relat_1 X1 (k1_rcomp_1 \\ & k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ & ((v5_relat_1 (k3_fuzzy_1 X0 X1) (k1_rcomp_1 k6_numbers np_1)) \wedge \\ & ((v1_funct_1 (k3_fuzzy_1 X0 X1)) \wedge ((v1_funct_2 (k3_fuzzy_1 X0 \\ & X1) X0 k1_numbers) \wedge (m1_subset_1 (k3_fuzzy_1 X0 X1) (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k1_numbers)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v5_relat_1 X1 (k1_rcomp_1 \\ k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ (\forall X2.((v5_relat_1 X2 (k1_rcomp_1 k6_numbers np_1)) \wedge \\ (v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 k1_numbers) \wedge (m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow ((r1_fuzzy_1 \\ X1 X2) \Leftrightarrow (r1_fuzzy_1 (k3_fuzzy_1 X0 X2) (k3_fuzzy_1 X0 X1)))) \end{aligned}$$