

t36_measure6 (TMQMAUaYVMPcSZ-
zGGecp9Z8GrDDgVhCdn6x)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_0 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow ((X0 = k6_numbers) \Leftrightarrow (k18_complex1 X0 = k6_numbers)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (4)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k7_real_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k18_complex1 X0 = k16_complex1 X0) \quad (8)$$

Assume the following.

$$\exists X0.(v1_xboole_0 X0) \wedge ((v1_xcmplx_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (v1_xreal_0 X0))) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow ((v1_xreal_0 (k16_complex1 X0)) \wedge (\neg v3_xxreal_0 (k16_complex1 X0))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(((v2_xxreal_0 X0) \wedge (v1_xreal_0 X0)) \wedge (\neg v3_xxreal_0 X1) \wedge (v1_xreal_0 X1)) \Rightarrow (v2_xxreal_0 (k2_xcmplx_0 X1 X0)) \quad (11)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k18_complex1 X0) k1_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (v1_xreal_0 (k16_complex1 X0)) \quad (13)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge ((\neg v2_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \Rightarrow ((v1_xboole_0 X0) \wedge (v1_xxreal_0 X0)) \quad (14)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0) \wedge (v2_xxreal_0 X0)) \Rightarrow ((\neg v1_xboole_0 X0) \wedge ((v1_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X0))) \quad (16)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (17)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k7_real_1 (k18_complex1 X0) (k18_complex1 X1) = k6_numbers) \Rightarrow (X0 = k6_numbers)))$$