

t36_subset_1

(TMajU5A3uE5QWts4HqmDugzebAARydywSHs)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.\neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.(X4 = k2_enumset1 X0 X1 X2 X3) \Leftrightarrow (\forall X5.(X5 \in X4) \Leftrightarrow (\neg(X5 \neq X0) \wedge ((X5 \neq X1) \wedge ((X5 \neq X2) \wedge (X5 \neq X3)))))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_zfmisc_1 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (r1_tarski X2 X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (X1 \in X0))) \wedge ((v1_xboole_0 X0) \Rightarrow ((m1_subset_1 X1 X0) \Leftrightarrow (v1_xboole_0 X1))) \quad (6)$$

Theorem 1

$$\forall X0.\forall X1.(m1_subset_1 X1 X0) \Rightarrow (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (\forall X3.(m1_subset_1 X3 X0) \Rightarrow (\forall X4.(m1_subset_1 X4 X0) \Rightarrow ((X0 \neq k1_xboole_0) \Rightarrow (m1_subset_1 (k2_enumset1 X1 X2 X3 X4) (k1_zfmisc_1 X0)))))))$$