

t370_xxreal_1
(TMGurfKq5rtLM6LvoakmKGLjMjDPzBqjyBw)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (r1_xxreal_0 X0 k1_xxreal_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X2 X1)) \Rightarrow \\ (k6_subset_1 (k1_xxreal_1 X0 X1) (k2_xxreal_1 X2 X1) = k2_xboole_0 \\ (k2_xxreal_1 X0 X2) (k1_tarski X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$v1_xxreal_0 k1_xxreal_0 \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (k6_subset_1 \\ (k1_xxreal_1 X0 k1_xxreal_0) (k2_xxreal_1 X1 k1_xxreal_0) = k2_xboole_0 \\ (k2_xxreal_1 X0 X1) (k1_tarski k1_xxreal_0))) \end{aligned}$$