

## t37\_lukasi\_1

(TMXQNAx2t5ak6BMjnwpciyhtFf3dZB8jRQn)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k3\_cqc\_lang : \iota \Rightarrow \iota$  be given. Let  $k8\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_cqc\_the1 : \iota \Rightarrow \iota$  be given. Let  $k6\_cqc\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X2.(m2\_subset\_1 X2 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (\forall X3.(m2\_subset\_1 X3 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (((k8\_cqc\_lang X0 X1 X2 \in k4\_cqc\_the1 X0) \wedge \\ (k8\_cqc\_lang X0 X2 X3 \in k4\_cqc\_the1 X0)) \Rightarrow (k8\_cqc\_lang X0 X1 X3 \in k4\_cqc\_the1 \\ X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (k8\_cqc\_lang X0 X1 (k6\_cqc\_lang X0 (k6\_cqc\_lang \\ X0 X1)) \in k4\_cqc\_the1 X0)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m2\_subset\_1 X1 (k9\_qc\_lang1 \\ X0) (k3\_cqc\_lang X0)) \Rightarrow (k8\_cqc\_lang X0 (k6\_cqc\_lang X0 (k6\_cqc\_lang \\ X0 X1)) X1 \in k4\_cqc\_the1 X0)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{4}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\neg v1\_xboole\_0 (k3\_cqc\_lang X0)) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((m1\_qc\_lang1\ X0)\wedge(m1\_subset\_1\ X1\ (k3\_cqc\_lang\ X0)))\Rightarrow(m2\_subset\_1\ (k6\_cqc\_lang\ X0\ X1)\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0)) \quad (6)$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1\ X0)\Rightarrow(m1\_subset\_1\ (k3\_cqc\_lang\ X0)\ (k1\_zfmisc\_1\ (k9\_qc\_lang1\ X0))) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1\ X0))\Rightarrow(v1\_xboole\_0\ X1)) \quad (8)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(m1\_qc\_lang1\ X0)\Rightarrow(\forall X1.(m2\_subset\_1\ X1\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0))\Rightarrow(\forall X2.(m2\_subset\_1\ X2\ (k9\_qc\_lang1\ X0)\ (k3\_cqc\_lang\ X0))\Rightarrow((k8\_cqc\_lang\ X0\ X1\ X2 \in k4\_cqc\_the1\ X0)\Leftrightarrow (k8\_cqc\_lang\ X0\ X1\ (k6\_cqc\_lang\ X0\ (k6\_cqc\_lang\ X0\ X2)) \in k4\_cqc\_the1\ X0)))) \end{aligned}$$