

t37_mesfunc5

(TMQ5nigCzq2TNmd8cqGjLSPNaPKS5KmkN9z)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_numbers : \iota$ be given. Let $v10_valued_0 : \iota \Rightarrow o$ be given. Let $v6_supinf_2 : \iota \Rightarrow o$ be given. Let $v4_measure1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_prob_2 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_tarski : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_supinf_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 \\ & X2)) \Rightarrow (((v1_prob_2 X2) \wedge (m2_finseq_1 X2 X1)) \Rightarrow (k3_tarski (k10_xtuple_0 \\ & X2) \in X1)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v5_relat_1 X1 X0))\Rightarrow(k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k7_numbers))))\Rightarrow((r1_mesfunc2 \\ X0 X1 X2)\Leftrightarrow((v3_valued_0 X2)\wedge(\exists X3.((v1_prob_2 X3)\wedge(m2_finseq_1 \\ X3 X1))\wedge((k1_relset_1 X0 X2 = k3_tarski (k2_relset_1 X1 X3))\wedge(\forall X4. \\ (v7_ordinal1 X4)\Rightarrow(\forall X5.(m1_subset_1 X5 X0)\Rightarrow(\forall X6. \\ (m1_subset_1 X6 X0)\Rightarrow(((X4 \in k4_finseq_1 X3)\wedge((X5 \in k1_funct_1 X3 \\ X4)\wedge(X6 \in k1_funct_1 X3 X4))\Rightarrow(k12_supinf_2 X2 X5 = k12_supinf_2 \\ X2 X6)))))))))) \quad (8) \end{aligned}$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (10)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((\neg v1_xboole_0 X1)\wedge \\ ((v1_prob_1 X1 X0)\wedge((v4_prob_1 X1 X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 \\ (k1_zfmisc_1 X0))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge((v1_funct_2 \\ X2 X1 k7_numbers)\wedge((v10_valued_0 X2)\wedge((v6_supinf_2 X2)\wedge((v4_measure1 \\ X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 k7_numbers))))))\Rightarrow \\ (\forall X3.((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 k7_numbers))))\Rightarrow((r1_mesfunc2 X0 X1 X3)\Rightarrow(m2_subset_1 (k1_relset_1 \\ X0 X3) (k1_zfmisc_1 X0 X1)))))) \quad (11) \end{aligned}$$