

t37_ordinal1

(TMJfS1nwrz2Su4DDHHA9JqCAXGiZgkPNoAN)

October 27, 2020

Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_ordinal1 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in k1_ordinal1 X1) \Leftrightarrow ((X0 \in X1) \vee (X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) (k1_tarski X1)) \Rightarrow (X0 = X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (X0 = X1) \Leftrightarrow ((r1_tarski X0 X1) \wedge (r1_tarski X1 X0)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (\neg X1 \in X0) \quad (7)$$

Theorem 1 $\forall X0. k6_subset_1 (k1_ordinal1 X0) (k1_tarski X0) = X0.$