

t37_quatern3
(TMYxccc3vfN93r8PLWF5SDAbp2gJhC6Y9b6a)

October 27, 2020

Let $v1_quaterni : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_quatern2 : \iota \Rightarrow \iota$ be given. Let $k29_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k32_quaterni : \iota \Rightarrow \iota$ be given. Let $k28_quaterni : \iota \Rightarrow \iota$ be given. Let $k26_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_quaterni : \iota \Rightarrow \iota$ be given. Let $k7_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k32_quaterni (k28_quaterni X0) = k32_quaterni X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (\forall X1.(v1_quaterni X1) \Rightarrow (\forall X2.(v1_quaterni X2) \Rightarrow (r1_xxreal_0 (k3_quatern2 (k29_quaterni (k26_quaterni X0 X1) X2)) (k7_real_1 (k7_real_1 (k3_quatern2 X0) (k3_quatern2 X1)) (k3_quatern2 X2)))))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k3_quatern2 X0 = k32_quaterni X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (k29_quaterni X0 X1 = k9_quaterni X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k28_quaterni X0 = k8_quaterni X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (k26_quaterni X0 X1 = k7_quaterni X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.(v1_quaterni\ X0) \Rightarrow (v1_quaterni\ (k8_quaterni\ X0)) \quad (7)$$

Assume the following.

$$\forall X0.(v1_quaterni\ X0) \Rightarrow (\forall X1.(v1_quaterni\ X1) \Rightarrow (k9_quaterni\ X0\ X1 = k7_quaterni\ X0\ (k8_quaterni\ X1))) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_quaterni\ X0) \Rightarrow (\forall X1.(v1_quaterni\ X1) \Rightarrow (\forall X2. \\ & (v1_quaterni\ X2) \Rightarrow (r1_xxreal_0\ (k3_quatern2\ (k29_quaterni\ (k29_quaterni \\ & X0\ X1)\ X2))\ (k7_real_1\ (k7_real_1\ (k3_quatern2\ X0)\ (k3_quatern2 \\ & X1))\ (k3_quatern2\ X2)))))) \end{aligned}$$