

t38_circcmb3

(TMc841s2z848yeZnexxToEWDFskBKUWDWnM)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_circcmb3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k3_msafree2 : \iota \Rightarrow \iota$ be given. Let $k3_circcmb3 : \iota \Rightarrow \iota$ be given. Let $k4_circcmb3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_msafree2 : \iota \Rightarrow \iota$ be given. Let $k6_circcmb3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\neg X0 \in X1) \Rightarrow (r1_xboole_0 (k1_tarski X0) X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1_xboole_0 X2) \wedge (v1_finset_1 \\ & X2)) \Rightarrow (\forall X3. (m1_subset_1 X3 k5_numbers) \Rightarrow (\forall X4. ((\\ & v1_funct_1 X4) \wedge ((v1_funct_2 X4 (k4_finseq_2 X3 X2) X2) \wedge (m1_subset_1 \\ & X4 (k1_zfmisc_1 (k2_zfmisc_1 (k4_finseq_2 X3 X2) X2)))))) \Rightarrow (\forall X5. \\ & ((v1_relat_1 X5) \wedge ((v1_funct_1 X5) \wedge ((v3_card_1 X5 X3) \wedge (v1_finseq_1 \\ & X5)))))) \Rightarrow (\forall X6. (m1_circcmb3 X6 X2) \Rightarrow (((k10_xtuple_0 X5 = k2_xboole_0 \\ & X0 X1) \wedge ((r1_tarski X0 (u1_struct_0 X6)) \wedge (r1_xboole_0 X1 (k3_msafree2 \\ & X6)))))) \Rightarrow ((k3_circcmb3 (k4_circcmb3 X3 X2 X4 X5) \in k2_msafree2 X6) \vee \\ & (k2_msafree2 (k6_circcmb3 X2 X6 (k4_circcmb3 X3 X2 X4 X5)) = k2_xboole_0 \\ & (k2_msafree2 X6) X1)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.k2_tarski\ X0\ X1 = k2_xboole_0\ (k1_tarski\ X0)\ (k1_tarski\ X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.k10_xtuple_0\ (k10_finseq_1\ X0\ X1) = k2_tarski\ X0\ X1 \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xreal_0\ np_2) \wedge (m2_subset_1\ np_2\ k1_numbers\ k5_numbers)) \wedge \\ & ((m1_subset_1\ np_2\ k5_numbers) \wedge (m1_subset_1\ np_2\ k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1\ (k10_finseq_1\ X0\ X1)) \wedge (v1_funct_1\ (k10_finseq_1\ X0\ X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v3_card_1\ (k10_finseq_1\ X0\ X1)\ np_2 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.v1_finseq_1\ (k10_finseq_1\ X0\ X1) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0\ X2) \wedge (v1_finset_1\ X2)) \Rightarrow (\forall X3.((v1_funct_1\ X3) \wedge ((v1_funct_2\ X3\ (k4_finseq_2\ np_2\ X2)\ X2) \wedge (m1_subset_1\ X3\ (k1_zfmisc_1\ (k2_zfmisc_1\ (k4_finseq_2\ np_2\ X2)\ X2)))))) \Rightarrow (\forall X4.(m1_circcmb3\ X4\ X2) \Rightarrow ((X0 \in u1_struct_0\ X4) \Rightarrow ((X1 \in k3_msafree2\ X4) \vee ((k3_circcmb3\ (k4_circcmb3\ np_2\ X2\ X3\ (k10_finseq_1\ X0\ X1)) \in k2_msafree2\ X4) \vee (k2_msafree2\ (k6_circcmb3\ X2\ X4\ (k4_circcmb3\ np_2\ X2\ X3\ (k10_finseq_1\ X0\ X1))) = k2_xboole_0\ (k2_msafree2\ X4)\ (k1_tarski\ X1))))))))) \end{aligned}$$