

t38\_finseq\_4  
(TMKGiD6Te22w4nF3Sdhieu8bVEGu7MW4Bxk)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_finseq\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1\_xboole\_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1)) \wedge (r1\_xboole\_0 X0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1. \neg(X1 \in k10\_xtuple\_0 X0) \wedge (X1 \in k10\_xtuple\_0 (k5\_finseq\_4 X0 X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (3)$$

**Theorem 1**

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1. (X1 \in k10\_xtuple\_0 X0) \Rightarrow (r1\_xboole\_0 (k10\_xtuple\_0 (k5\_finseq\_4 X0 X1)) (k1\_tarski X1)))$$