

t38_funct_2
(TMZQ5jSNPAuX4esiuYjyomPyq4TuKfkCPJ9)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k8_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k8_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k8_relat_1 X2 X1) \Leftrightarrow ((k4_tarski X0 (k1_funct_1 X2 X0) \in X2) \wedge (k1_funct_1 X2 X0 \in X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k4_tarski X0 X1 \in X2) \Leftrightarrow ((X0 \in k9_xtuple_0 X2) \wedge (X1 = k1_funct_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k8_relset_1 X0 X1 X2 X3 = k8_relat_1 X2 X3) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (((X1 \neq k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X0 = k1_relset_1 X0 X2))) \wedge ((X1 = k1_xboole_0) \Rightarrow ((v1_funct_2 X2 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (7)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X3)\wedge((v1_funct_2 X3 X0 X1)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))\Rightarrow((X1\neq k1_xboole_0)\Rightarrow(\forall X4.(X4 \in k8_relset_1 X0 X1 X3 X2)\Leftrightarrow((X4 \in X0)\wedge(k1_funct_1 X3 X4 \in X2))))$$