

t38\_funct\_4 (TMZWvjY-  
HjzDu6Xa9Ev4jJTEFKZdbqM6ZKZg)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0. \forall X1. \forall X2. ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\
 & X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow \\
 & (\forall X3. ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\
 & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((X1 = k1\_xboole\_0) \Rightarrow \\
 & (X0 = k1\_xboole\_0)) \Rightarrow (k1\_funct\_4 X2 X3 = X3)))
 \end{aligned} \tag{1}$$

**Theorem 1**

$$\begin{aligned}
 & \forall X0. \forall X1. ((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 X0 X0) \wedge \\
 & (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0)))) \Rightarrow (\forall X2. \\
 & ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 X0 X0) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\
 & (k2\_zfmisc\_1 X0 X0)))) \Rightarrow (k1\_funct\_4 X1 X2 = X2))
 \end{aligned}$$