

t38_topgen_3
(TMcppk5h9GzNS4XjZedYJvs5PaWVy8SZLB9)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_topgen_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v1_pre_topc : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. r1_tarski X0 (k2_xboole_0 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ (\forall X2. ((\neg v1_xboole_0 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (u1_struct_0 (k6_topgen_3 X0 X1)))))) \Rightarrow (k2_pre_topc (k6_topgen_3 \\ X0 X1) X2 = k2_xboole_0 X2 (k6_domain_1 X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ (u1_struct_0 X0))) \Rightarrow (((v4_pre_topc X1 X0) \Rightarrow (k2_pre_topc X0 X1 = \\ X1)) \wedge (((v2_pre_topc X0) \wedge (k2_pre_topc X0 X1 = X1)) \Rightarrow (v4_pre_topc \\ X1 X0)))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k2_xboole_0 X0 X1 = X1) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow \\ (k6_domain_1 X0 X1 = k1_tarski X1) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_pre_topc (k6_topgen_3 X0 X1))\wedge((v2_pre_topc (k6_topgen_3 X0 X1))\wedge(l1_pre_topc (k6_topgen_3 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (8)$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow \\ (\forall X2.((\neg v1_xboole_0 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 \\ (u1_struct_0 (k6_topgen_3 X0 X1))))))\Rightarrow((v4_pre_topc X2 (k6_topgen_3 \\ X0 X1))\Leftrightarrow(X1 \in X2)))) \end{aligned}$$