

t390\_xxreal\_1

(TMHtJGA7nvTgnSRoQwuE7pkYvy58ctKR5rm)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k6\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xxreal\_0 : \iota$  be given. Let  $k1\_xxreal\_0 : \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k2\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xxreal\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (r1\_xxreal\_0 X0 k1\_xxreal\_0) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 \\ X1 X0) \Rightarrow (k6\_subset\_1 (k3\_xxreal\_1 k2\_xxreal\_0 X0) (k1\_tarski X1) = \\ k2\_xboole\_0 (k4\_xxreal\_1 k2\_xxreal\_0 X1) (k3\_xxreal\_1 X1 X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$v1\_xxreal\_0 k1\_xxreal\_0 \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (4)$$

**Theorem 1**

$$\begin{aligned} \forall X0.(v1\_xreal\_0 X0) \Rightarrow (k6\_subset\_1 (k3\_xxreal\_1 k2\_xxreal\_0 \\ k1\_xxreal\_0) (k1\_tarski X0) = k2\_xboole\_0 (k4\_xxreal\_1 k2\_xxreal\_0 \\ X0) (k3\_xxreal\_1 X0 k1\_xxreal\_0)) \end{aligned}$$