

t39_bvfunc11

(TMU6v8fLV1ZqVK7gSqTBSHVKKQXUJ7wg9nXS)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_2 : \iota \Rightarrow \iota$ be given. Let $m1_eqrel_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v2_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_bvfunc_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_partit1 : \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_margrel1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ & (k1_zfmisc_1 (k1_bvfunc_2 X0))) \Rightarrow (\forall X3. (m1_eqrel_1 X3 X0) \Rightarrow \\ & (\forall X4. (m1_eqrel_1 X4 X0) \Rightarrow ((v2_bvfunc_2 X2 X0) \Rightarrow (r1_bvfunc_1 \\ & X0 (k7_bvfunc_2 X0 (k1_bvfunc_1 X0 (k7_bvfunc_2 X0 X1 X2 X3)) X2 X4) \\ & (k1_bvfunc_1 X0 (k6_bvfunc_2 X0 (k6_bvfunc_2 X0 X1 X2 X4) X2 X3))))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k1_partit1 X0))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 \\ & X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 k6_margrel1)))) \Rightarrow (\forall X3. (m1_eqrel_1 X3 X0) \Rightarrow (r2_funct_2 \\ & X0 k6_margrel1 (k1_bvfunc_1 X0 (k6_bvfunc_2 X0 X2 X1 X3)) (k7_bvfunc_2 \\ & X0 (k1_bvfunc_1 X0 X2) X1 X3)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \wedge ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((r2_funct_2 X0 X1 X2 \\ & X3) \Leftrightarrow (X2 = X3)) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.k1.bvfunc.2 X0 = k1.partit1 X0 \quad (4)$$

Assume the following.

$$\begin{aligned} &\forall X0.\forall X1.((\neg v1.xboole.0 X0)\wedge((v1.funct.1 X1)\wedge \\ &(v1.funct.2 X1 X0 k6.margrel1)\wedge(m1.subset.1 X1 (k1.zfmisc.1 (\\ &k2.zfmisc.1 X0 k6.margrel1))))))\Rightarrow(k1.bvfunc.1 X0 X1 = k13.margrel1 \\ &X1) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} &\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1.xboole.0 X0)\wedge \\ &(((v1.funct.1 X1)\wedge((v1.funct.2 X1 X0 k6.margrel1)\wedge(m1.subset.1 \\ &X1 (k1.zfmisc.1 (k2.zfmisc.1 X0 k6.margrel1))))))\wedge((m1.subset.1 \\ &X2 (k1.zfmisc.1 (k1.bvfunc.2 X0)))\wedge(m1.eqrel.1 X3 X0))))\Rightarrow((v1.funct.1 \\ &(k7.bvfunc.2 X0 X1 X2 X3)\wedge((v1.funct.2 (k7.bvfunc.2 X0 X1 X2 X3) \\ &X0 k6.margrel1)\wedge(m1.subset.1 (k7.bvfunc.2 X0 X1 X2 X3) (k1.zfmisc.1 \\ &(k2.zfmisc.1 X0 k6.margrel1)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} &\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1.xboole.0 X0)\wedge \\ &(((v1.funct.1 X1)\wedge((v1.funct.2 X1 X0 k6.margrel1)\wedge(m1.subset.1 \\ &X1 (k1.zfmisc.1 (k2.zfmisc.1 X0 k6.margrel1))))))\wedge((m1.subset.1 \\ &X2 (k1.zfmisc.1 (k1.bvfunc.2 X0)))\wedge(m1.eqrel.1 X3 X0))))\Rightarrow((v1.funct.1 \\ &(k6.bvfunc.2 X0 X1 X2 X3)\wedge((v1.funct.2 (k6.bvfunc.2 X0 X1 X2 X3) \\ &X0 k6.margrel1)\wedge(m1.subset.1 (k6.bvfunc.2 X0 X1 X2 X3) (k1.zfmisc.1 \\ &(k2.zfmisc.1 X0 k6.margrel1)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} &\forall X0.\forall X1.((\neg v1.xboole.0 X0)\wedge((v1.funct.1 X1)\wedge \\ &(v1.funct.2 X1 X0 k6.margrel1)\wedge(m1.subset.1 X1 (k1.zfmisc.1 (\\ &k2.zfmisc.1 X0 k6.margrel1))))))\Rightarrow((v1.funct.1 (k1.bvfunc.1 \\ &X0 X1))\wedge((v1.funct.2 (k1.bvfunc.1 X0 X1) X0 k6.margrel1)\wedge(m1.subset.1 \\ &(k1.bvfunc.1 X0 X1) (k1.zfmisc.1 (k2.zfmisc.1 X0 k6.margrel1)))))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} &\forall X0.(\neg v1.xboole.0 X0)\Rightarrow(\forall X1.((v1.funct.1 X1)\wedge \\ &(v1.funct.2 X1 X0 k6.margrel1)\wedge(m1.subset.1 X1 (k1.zfmisc.1 (\\ &k2.zfmisc.1 X0 k6.margrel1))))))\Rightarrow(\forall X2.(m1.subset.1 X2 \\ &(k1.zfmisc.1 (k1.bvfunc.2 X0)))\Rightarrow(\forall X3.(m1.eqrel.1 X3 X0)\Rightarrow \\ &(\forall X4.(m1.eqrel.1 X4 X0)\Rightarrow((v2.bvfunc.2 X2 X0)\Rightarrow(r1.bvfunc.1 \\ &X0 (k1.bvfunc.1 X0 (k6.bvfunc.2 X0 (k7.bvfunc.2 X0 X1 X2 X3) X2 X4)) \\ &(k7.bvfunc.2 X0 (k1.bvfunc.1 X0 (k6.bvfunc.2 X0 X1 X2 X4) X2 X3)))))) \end{aligned}$$