

t39_bvfunc25

(TMEg4yK5RP6kSSh8FouVacqC3QWbrT5NgFJ)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X3.((v1_funct_1 X3) \wedge \\
 & ((v1_funct_2 X3 X0 k6_margrel1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((r2_funct_2 X0 k6_margrel1 \\
 & (k9_bvfunc_1 X0 X1 (k9_bvfunc_1 X0 X2 X3)) (k12_bvfunc_1 X0)) \Rightarrow (\\
 & r2_funct_2 X0 k6_margrel1 (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 X1 X2) \\
 & (k9_bvfunc_1 X0 X1 X3)) (k12_bvfunc_1 X0))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (\\
 & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\
 & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\
 & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\
 & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\
 & (k9_bvfunc_1 X0 X2 (k9_bvfunc_1 X0 X1 X2)) (k12_bvfunc_1 X0))))
 \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ (v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 X1 X2) (k9_bvfunc_1 X0 X1 X1)) (k12_bvfunc_1 \\ X0))) \end{aligned}$$