

t39_finseq_5
(TMHNKg3ySCRESkL3hCPswkY2SUdiGQLigjX)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & \quad (\forall X1. \forall X2. (v7_ordinal1 X2) \Rightarrow (\neg(X2 \in k4_finseq_1 X0) \wedge \\ & \quad ((\neg r1_xxreal_0 (k4_finseq_4 X0 X1) X2) \wedge (k1_funct_1 X0 X2 = X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & \quad (k4_finseq_1 X0 = k9_xtuple_0 X0) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & \quad (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & \quad X0)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v5_relat_1 X1 X0) \wedge (\\ & \quad v1_funct_1 X1))) \Rightarrow (\forall X2. (X2 \in k9_xtuple_0 X1) \Rightarrow (k7_partfun1 \\ & \quad X0 X1 X2 = k1_funct_1 X1 X2)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (7)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(m2_finseq_1 X2 X1)\Rightarrow((X0 \in k4_finseq_1 X2)\Rightarrow(r1_xxreal_0 (k4_finseq_4 X2 (k7_partfun1 X1 X2 X0)) X0))))$$