

t39_glib_003 (TMTbtEPVbd- dLvzYH9GE9ApM6si9gBQrkyUd)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $v3_glib_003 : \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_003 : \iota \Rightarrow \iota$ be given. Let $k13_glib_003 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. k1_funct_1 (k16_funcop_1 X0 X1) X0 = X1 \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_003 X0)))))) \Rightarrow \\ & (\forall X1. \forall X2. (X1 \in k6_glib_000 X0) \Rightarrow (k7_glib_003 (k13_glib_003 \\ & X0 X1 X2) = k1_funct_4 (k7_glib_003 X0) (k16_funcop_1 X1 X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 \\ & (k1_funct_4 X2 X1) X0 = k1_funct_1 X1 X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 (k2_funcop_1 X0 X1))\wedge(v1_funct_1 (k2_funcop_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_003 X0))))))\Rightarrow ((v1_relat_1 (k7_glib_003 X0))\wedge(v1_funct_1 (k7_glib_003 X0))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow (X2 = X0)) \quad (9)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_003 X0))))))\Rightarrow (\forall X1.\forall X2.(X1 \in k6_glib_000 X0)\Rightarrow(k1_funct_1 (k7_glib_003 (k13_glib_003 X0 X1 X2)) X1 = X2))$$