

t39_int_5

(TML5fEV9rygLY6Gig22PpmMS3XUtBjzjKB2)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k2_finseq_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & ((v2_funct_1 X0) \Leftrightarrow (k5_card_1 (k10_xtuple_0 X0) = k3_finseq_1 X0)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 X0) \Rightarrow \\ & (\forall X2.(v7_ordinal1 X2) \Rightarrow (\forall X3.(v7_ordinal1 X3) \Rightarrow (\\ & k10_xtuple_0 (k2_finseq_7 X0 X1 X2 X3) = k10_xtuple_0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m2_finseq_1 X1 X0) \Rightarrow \\ & (\forall X2.(v7_ordinal1 X2) \Rightarrow (\forall X3.(v7_ordinal1 X3) \Rightarrow (\\ & k3_finseq_1 (k2_finseq_7 X0 X1 X2 X3) = k3_finseq_1 X1)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (\\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge(v1_funct_1 X1)\wedge(v1_finseq_1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge((m1_finseq_1 X1 X0)\wedge((v7_ordinal1 X2)\wedge(v7_ordinal1 X3))))\Rightarrow(m2_finseq_1 (k2_finseq_7 X0 X1 X2 X3) X0) \quad (7)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow(\forall X3.(v7_ordinal1 X3)\Rightarrow((v2_funct_1 X1)\Leftrightarrow(v2_funct_1 (k2_finseq_7 X0 X1 X2 X3))))))$$