

t39\_xreal\_1  
(TMP5hwtcdZfUNZjb9gytfMdvaA6EG5gxApc)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 k6\_numbers) \wedge (r1\_xxreal\_0 (k2\_xcmplx\_0 X1 X0) X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (v1\_xreal\_0 (k2\_xcmplx\_0 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k2\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (5)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 k6\_numbers) \wedge ((r1\_xxreal\_0 X1 X2) \wedge (r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X2) X1))))))$$