

t3_afinsq_2
(TMHSX2RgLU1ySuGARmZXcLfVY5P5QNwywv8)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v5_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (4)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (5)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.(X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.(X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0)))) \Rightarrow ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v5_ordinal1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0))))) \quad (9)$$

Theorem 1

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge ((v5_ordinal1 X1) \wedge (v1_finset_1 X1)))) \Rightarrow (\neg(X0 \in k10_xtuple_0 X1) \wedge (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (\neg(X2 \in k1_relset_1 k5_numbers X1) \wedge (k1_funct_1 X1 X2 = X0))))$$