

t3_arytm_3

(TMKUP8SyUpYjTuAyTeLgUeerbaufVELvf2x)

October 27, 2020

Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_arytm_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $np_1 : \iota$ be given. Let $k11_ordinal2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_ordinal1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow ((k11_ordinal2 np_1 X0 = X0) \wedge (k11_ordinal2 X0 np_1 = X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow (k11_ordinal2 X0 k1_xboole_0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$np_1 = k1_ordinal1 k1_xboole_0 \quad (3)$$

Assume the following.

$$\forall X0.(v3_ordinal1 X0) \Rightarrow ((\neg v1_xboole_0 (k1_ordinal1 X0)) \wedge (v3_ordinal1 (k1_ordinal1 X0))) \quad (4)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.(v3_ordinal1 X0) \Rightarrow (\forall X1.(v3_ordinal1 X1) \Rightarrow ((\\ r1_arytm_3 X0 X1) \Leftrightarrow (\forall X2.(v3_ordinal1 X2) \Rightarrow (\forall X3.(\\ v3_ordinal1 X3) \Rightarrow (\forall X4.(v3_ordinal1 X4) \Rightarrow (((X0 = k11_ordinal2 \\ X2 X3) \wedge (X1 = k11_ordinal2 X2 X4)) \Rightarrow (X2 = np_1))))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (v3_ordinal1 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v3_ordinal1 X0) \Rightarrow ((r1_arytm_3 k1_xboole_0 X0) \Rightarrow (X0 = np_1))$$