

t3_cfunct_1

(TMRQhnt5airDzpZf4c4gMsH1mys8pzBmyW)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k19_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_complex1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $k18_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k2_numbers) \wedge (m1_subset_1 X1 k2_numbers)) \Rightarrow (k9_complex1 X0 X1 = k3_xcmplx_0 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. \forall X4. ((v1_membered X1) \wedge ((v1_membered X2) \wedge (((v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \wedge ((v1_funct_1 X4) \wedge (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2))))))) \Rightarrow (k19_valued_1 X0 X1 X2 X3 X4 = k18_valued_1 X3 X4) \quad (4)$$

Assume the following.

$$v1_membered k2_numbers \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(m1_subset_1 (k7_partfun1 X0 X1 X2) X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v1_membered X1)\wedge((v1_membered X2)\wedge(((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge((v1_funct_1 X4)\wedge(m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2)))))))\Rightarrow((v1_funct_1 (k19_valued_1 X0 X1 X2 X3 X4))\wedge(m1_subset_1 (k19_valued_1 X0 X1 X2 X3 X4) (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_valued_0 X1))))\Rightarrow((v1_relat_1 (k18_valued_1 X0 X1))\wedge(v1_funct_1 (k18_valued_1 X0 X1))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k3_xboole_0 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in X0)\wedge(X3 \in X1))) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_valued_0 X1)))\Rightarrow(\forall X2.((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((X2 = k18_valued_1 X0 X1)\Leftrightarrow((k9_xtuple_0 X2 = k3_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1))\wedge(\forall X3.(X3 \in k9_xtuple_0 X2)\Rightarrow(k1_funct_1 X2 X3 = k3_xcmplx_0 (k1_funct_1 X0 X3) (k1_funct_1 X1 X3))))))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_valued_0 X2)) \quad (15)$$

Theorem 1

$$\begin{aligned} &\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge(\\ &\quad m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers))))\Rightarrow(\\ &\quad \forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ &\quad X0 k2_numbers))))\Rightarrow((k1_relset_1 X0 (k19_valued_1 X0 k2_numbers \\ &\quad k2_numbers X1 X2) = k9_subset_1 X0 (k1_relset_1 X0 X1) (k1_relset_1 \\ &\quad X0 X2))\wedge(\forall X3.(m1_subset_1 X3 X0)\Rightarrow((X3 \in k1_relset_1 X0 (\\ &\quad k19_valued_1 X0 k2_numbers k2_numbers X1 X2))\Rightarrow(k7_partfun1 k2_numbers \\ &\quad (k19_valued_1 X0 k2_numbers k2_numbers X1 X2) X3 = k9_complex1 (\\ &\quad k7_partfun1 k2_numbers X1 X3) (k7_partfun1 k2_numbers X2 X3)))))) \end{aligned}$$