

# t3\_e\_siec (TMLsYcQB- SNYZ8wmtvKUkBFvozvoPZQenqy1)

October 27, 2020

Let  $v2\_e\_siec : \iota \Rightarrow o$  be given. Let  $g1\_e\_siec : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $v3\_e\_siec : \iota \Rightarrow o$  be given. Let  $l1\_e\_siec : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k3\_relat\_1 (k4\_relat\_1 X0) X2 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (k4\_xboole\_0 X0 X1 = k1\_xboole\_0) \Leftrightarrow (r1\_tarski X0 X1) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (v1\_relat\_1 X1) \Rightarrow (\forall X2. (v1\_relat\_1 \\ & X2) \Rightarrow (((v2\_e\_siec (g1\_e\_siec X0 X1 X2)) \wedge ((v3\_e\_siec (g1\_e\_siec \\ & X0 X1 X2)) \wedge (l1\_e\_siec (g1\_e\_siec X0 X1 X2)))) \Leftrightarrow ((r1\_tarski X1 (k2\_zfmisc\_1 \\ & X0 X0)) \wedge ((r1\_tarski X2 (k2\_zfmisc\_1 X0 X0)) \wedge ((k3\_relat\_1 X1 X1 = \\ & X1) \wedge ((k3\_relat\_1 X1 X2 = X1) \wedge ((k3\_relat\_1 X2 X2 = X2) \wedge ((k3\_relat\_1 \\ & X2 X1 = X2) \wedge ((k3\_relat\_1 X1 (k4\_xboole\_0 X1 (k4\_relat\_1 X0)) = k1\_xboole\_0) \wedge \\ & (k3\_relat\_1 X2 (k4\_xboole\_0 X2 (k4\_relat\_1 X0)) = k1\_xboole\_0)))))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. k3\_relat\_1 (k4\_relat\_1 X0) (k4\_relat\_1 X0) = k4\_relat\_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. m1\_subset\_1 k1\_xboole\_0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.r1\_tarSKI X0 X0 \quad (6)$$

Assume the following.

$$\forall X0.r1\_tarSKI (k4\_relat\_1 X0) (k2\_zfmisc\_1 X0 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k4\_relat\_1 X0))\wedge(v1\_funct\_1 (k4\_relat\_1 X0)) \quad (8)$$

**Theorem 1**

$$\forall X0.(v2\_e\_siec (g1\_e\_siec X0 (k4\_relat\_1 X0) (k4\_relat\_1 X0)))\wedge((v3\_e\_siec (g1\_e\_siec X0 (k4\_relat\_1 X0) (k4\_relat\_1 X0)))\wedge(l1\_e\_siec (g1\_e\_siec X0 (k4\_relat\_1 X0) (k4\_relat\_1 X0))))$$