

t3_funct_6

(TMY5dbdPZv4MVkPb8P5yXRKLsHL6mfFPHLD)

October 27, 2020

Let $k2_funct_4 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_binop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (X1 \in X0) \Rightarrow (k1_funct_1 (k2_funcop_1 X0 X2) X1 = X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((k9_xtuple_0 X2 = k2_zfmisc_1 X0 X1) \Rightarrow (k9_xtuple_0 (k2_funct_4 X2) = k2_zfmisc_1 X1 X0)) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((k9_xtuple_0 X0 = k9_xtuple_0 X1) \wedge (\forall X2. (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2))) \Rightarrow (X0 = X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 (k2_funcop_1 X0 X1))\wedge(v1_funct_1 (k2_funcop_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1))\wedge((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarski X1))\wedge(m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarski X1)))))) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow((v1_relat_1 (k2_funct_4 X0))\wedge(v1_funct_1 (k2_funct_4 X0))) \quad (8)$$

Assume the following.

$$\forall X0.(((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(((v1_relat_1 X1)\wedge(v1_funct_1 X1))\Rightarrow((X1 = k2_funct_4 X0)\Leftrightarrow((\forall X2.(X2 \in k9_xtuple_0 X1)\Leftrightarrow(\exists X3.\exists X4.(X2 = k4_tarski X4 X3)\wedge(k4_tarski X3 X4 \in k9_xtuple_0 X0))))\wedge(\forall X2.\forall X3.(k4_tarski X2 X3 \in k9_xtuple_0 X0)\Rightarrow(k1_binop_1 X1 X3 X2 = k1_binop_1 X0 X2 X3)))))) \quad (9)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.k1_binop_1 X0 X1 X2 = k1_funct_1 X0 (k4_tarski X1 X2)) \quad (10)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.k2_funct_4 (k7_funcop_1 (k2_zfmisc_1 X0 X1) X2) = k7_funcop_1 (k2_zfmisc_1 X1 X0) X2$$