

t3_group_11

(TMR6k5MGkBJ3T9KAECehJX4TtG9jbxGCmW1)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_group_1 : \iota \Rightarrow o$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_group_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k13_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_group_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_struct_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1)) \wedge (r1_xboole_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge (l3_algstr_0 X0))) \Rightarrow (\forall X1. (m1_group_2 X1 X0) \Rightarrow ((\neg v2_struct_0 X1) \wedge ((v2_group_1 X1) \wedge (l3_algstr_0 X1)))) \quad (3)$$

Assume the following.

$$\forall X0. (l3_algstr_0 X0) \Rightarrow (l1_struct_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \wedge (m1_group_2 X1 X0)) \Rightarrow (m1_subset_1 (k8_group_2 X0 X1) (k1_zfmisc_1 (u1_struct_0 X0))) \quad (5)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (k8_group_2 X0 X1 = u1_struct_0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(l1_struct_0 X0) \Rightarrow (\forall X1.(r1_struct_0 X0 X1) \Leftrightarrow (X1 \in u1_struct_0 X0)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_group_1 X0) \wedge ((v3_group_1 X0) \wedge (l3_algstr_0 X0)))) \Rightarrow (\forall X1.(m1_group_2 X1 X0) \Rightarrow (\forall X2. \\ & (m1_group_2 X2 X0) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow \\ & (\neg(\neg r1_xboole_0 (k13_group_2 X0 X1 X3) (k8_group_2 X0 X2)) \wedge (\forall X4. \\ & (m1_subset_1 X4 (u1_struct_0 X0)) \Rightarrow (\neg(X4 \in k13_group_2 X0 X1 X3) \wedge \\ & (r1_struct_0 X2 X4))))))) \end{aligned}$$