

t3\_jordan5b (TMJVfg-  
KEee1cE9U9dmJN5XVnKUeV7LBH6bM)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_nat\_d : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v6\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k7\_nat\_d X1 X2)) \wedge (r1\_xxreal\_0 \\ & X2 X1)) \Rightarrow (r1\_xxreal\_0 (k2\_xcmplx\_0 X0 X2) X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Rightarrow (r1\_xxreal\_0 (k7\_nat\_d \\ & X2 X1) (k7\_nat\_d X2 X0)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (k7\_nat\_d \\ & (k2\_xcmplx\_0 X0 X1) X1 = X0)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 \\ & X1)) \Rightarrow (k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \end{aligned} \quad (6)$$

Assume the following.

$$v6\_membered\ k4\_ordinal1 \tag{7}$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1\ X0)\wedge(v7\_ordinal1\ X1))\Rightarrow(v7\_ordinal1\ (k2\_xcmplx\_0\ X0\ X1)) \tag{8}$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1\ X0)\wedge(v7\_ordinal1\ X1))\Rightarrow(m1\_subset\_1\ (k7\_nat\_d\ X0\ X1)\ k5\_numbers) \tag{9}$$

Assume the following.

$$\forall X0.(v6\_membered\ X0)\Rightarrow(\forall X1.(m1\_subset\_1\ X1\ X0)\Rightarrow(v7\_ordinal1\ X1)) \tag{10}$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1\ X0)\Rightarrow(\forall X1.(v7\_ordinal1\ X1)\Rightarrow((r1\_xxreal\_0\ np\_1\ X0)\wedge(r1\_xxreal\_0\ np\_1\ X1))\Rightarrow(r1\_xxreal\_0\ (k2\_nat\_1\ (k7\_nat\_d\ X1\ X0)\ np\_1)\ X1)))$$