

t3_matrix_1

(TMFXoxSzJsQacveexwzGJSbKuzjdKrg1uis)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v1_matrix_1 : \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k4_finseq_1 X1 = k2_finseq_1 np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (1)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (3)$$

Assume the following.

$$\forall X0. v1_finseq_1 (k5_finseq_1 X0) \quad (4)$$

Assume the following.

$$\forall X0. (v1_relat_1 (k5_finseq_1 X0)) \wedge (v1_funct_1 (k5_finseq_1 X0)) \quad (5)$$

Assume the following.

$$\forall X0. (v1_finset_1 X0) \Rightarrow ((v1_finset_1 (k1_card_1 X0)) \wedge (v1_card_1 (k1_card_1 X0))) \quad (6)$$

Assume the following.

$$\forall X0.v1_card_1 (k1_card_1 X0) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1_tarski X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ ((v1_matrix_1 X0) \Leftrightarrow (\exists X1.(v7_ordinal1 X1) \wedge (\forall X2. \\ \neg(X2 \in k10_xtuple_0 X0) \wedge (\forall X3.((v1_relat_1 X3) \wedge ((v1_funct_1 \\ X3) \wedge (v1_finseq_1 X3)))) \Rightarrow (\neg(X3 = X2) \wedge (k3_finseq_1 X3 = X1)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.((v3_ordinal1 X0) \wedge (v1_finset_1 X0)) \Rightarrow (v7_ordinal1 X0) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finset_1 X0))) \quad (11)$$

Assume the following.

$$\forall X0.(v1_card_1 X0) \Rightarrow (v3_ordinal1 X0) \quad (12)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (v1_matrix_1 (k9_finseq_1 X0))$$