

t3_menelaus (TMWsRF- SJXp2wZLwQX4TLwLwtXGCd6avtePk)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k4_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_binop_2 : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (k4_algstr_0 (k15_euclid np_2) X0 = k19_euclid (k1_real_1 (k17_euclid \\ & X0)) (k1_real_1 (k18_euclid X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid \\ & (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (k7_binop_2 X0 = k4_xcmplx_0 X0) \tag{3}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k1_real_1 X0 = k4_xcmplx_0 X0) \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (m1_subset_1 (k1_real_1 \\ & X0) k1_numbers) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (m1_subset_1 (k18_euclid X0) k1_numbers) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (m1_subset_1 (k17_euclid X0) k1_numbers) \end{aligned} \tag{7}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((k17_euclid (k4_algstr_0 (k15_euclid np_2) X0) = k7_binop_2 \\ & (k17_euclid X0)) \wedge (k18_euclid (k4_algstr_0 (k15_euclid np_2) \\ & X0) = k7_binop_2 (k18_euclid X0))) \end{aligned}$$